arXiv:1003.2264v1 [quant-ph] 11 Mar 2010

Coherent States for PT-/Non-PT-Symmetric and Non-Hermitian Morse Potential via Path Integral Method

Nalan Kandirmaz[1,*] and Ramazan Sever[2,†]

[1]*Department of Physics, Mersin University, Mersin,Turkey*

[2]*Department of Physics, Middle East Technical University, Ankara,Turkey*

(Dated: March 12, 2010)

Abstract

We discuss the coherent states for PT-/non-PT-Symmetric and non-Hermitian generalized Morse Potential obtained by using path integral formalism over the holomorphic coordinates. We transform the action of generalized Morse potential into two harmonic oscillators with a new parametric time to establish the parametric time coherent states. We calculate the energy eigenvalues and the corresponding wave functions in parabolic coordinates.

Keywords: PT-/non-PT-symmetry, non-Hermitian, coherent states, path integral, Morse potential

PACS numbers: 31.15.Kb, 11.30Er

---

[*] E-mail: `nkandirmaz@mersin.edu.tr`

[†] E-mail: `sever@metu.edu.tr`

I. INTRODUCTION

PT-symmetric quantum mechanics has been widely investigated recently. In standard formalism of quantum mechanics, Hamiltonian that defines the dynamics and symmetries of system is Hermitian. In the PT-symmetric case, Hamiltonian has real spectra although it is not Hermitian. Bender and his co-workers studied PT-symmetric Hamiltonians which have real eigenvalues [1]. Following these works, many authors studied PT-symmetric and non-Hermitian Hamiltonians having real and/or complex eigenvalues [2, 3, 5] by applying the analytic methods and numerical techniques. In this direction, several methods have been developed to calculate energy eigenvalues and the wave functions [4]. One of the methods providing exact solutions is Path integral method . It is also employed to find the transition amplitudes among the configuration space eigenstates [6]. The energy spectra and eigenfunctions of PT-/non-PT- and non-Hermitian Morse potential is obteined [7].

Morse potential which has an important role in molecular physics is used for the description of the interaction between the atoms in diatomic molecules [8–10]. Bound-state energy spectrum and the wave functions of Morse potentials is obtained by Duru using path integral method [11]. Later Unal derived parametric time coherent states using path integrals over holomorphic coordinates [9]. Coherent states which are firstly constructed by Schrödinger are defined as eigenstates of lowering operators for harmonic oscillators [14]. They are minimum uncertainty states and do not disperse during the evolution of the system [9, 15, 16]. Both PT-symmetric and non-Hermitian generalized Morse potential have been studied exactly [7, 13]. In the present work coherent states of PT-/non-PT- and non-Hermitian Morse potential is discussed by using Path Integral method. We obtain the classical dynamics in terms of the holomorphic coordinates because of that holomorphic coordinates are the classical analogue of the rising and lowering operators of the harmonic oscillators. Energy eigenvalues and the corresponding wave functions are obtained by transforming the problem into the two oscillators in the parametric time and quantizing these oscillators using the path integration over the holomorphic coordinates. The path integral technique can be applied to any potential if it is transformed into the Harmonic oscillator potential.

The organization of the paper is as follows: In Sec. 2, we introduce coherent states of the generalized Morse potential. We obtain the energy eigenvalues and wave functions in parabolic coordinates. In Secs. 3 and 4, we get the coherent states, energy eigenvalues and corresponding eigenfunctions of the PT-symmetric and non-PT-symmetric non-Hermitian forms of the generalized Morse potential. Finally, Results are discussed in Sec. 5.

II. GENERALIZED MORSE POTENTIAL

We consider a particle with mass m moving in generalized Morse potential

$$V(x) = V_1 e^{-2\alpha x} - V_2 e^{-\alpha x} \tag{1}$$

where V_1, V_2 are constants and α is the width of the potential. The action of the one dimensional generalized Morse potential is

$$A = \int dt \left[p_x \frac{dx}{dt} - \left(\frac{p_x^2}{2m} + V_1 e^{-2\alpha x} - V_2 e^{-\alpha x} \right) \right]. \tag{2}$$

Let us define a new coordinate u(t)

$$x = -2\alpha \ln u \tag{3}$$

and canonical conjugate momentum

$$p_x = -\frac{\alpha u}{2} p_u. \tag{4}$$

After this transformation, action takes

$$A = \int dt \left[p_u \dot{u} - \left(\frac{\alpha^2 u^2}{4} \frac{p_u^2}{2m} + V_1 u^4 - V_2 u^2 \right) \right]. \tag{5}$$

In order to annihilate the factorization u^2 in kinetic energy term, we define a new time parameter as

$$\frac{dt}{d\tau} = \frac{1}{u^2} \tag{6}$$

Substitution of this new time parameter into Eq. (5) gives us

$$A = \int d\tau \left[p_u \frac{du}{d\tau} - \left(\frac{\alpha^2}{4} \frac{p_u^2}{2m} + V_1 u^2 - V_2 \right) + (-p_0) \left(\frac{dt}{d\tau} - \frac{1}{u^2} \right) \right]. \tag{7}$$

Here an additional Lagrange multiplier p_0 is used due to the presence of a new parametric time. Adding a dummy momentum p_ϕ by $\frac{d\phi}{d\tau}$ another Lagrange multiplier. Action becomes

$$A = \int d\tau \left[p_u \frac{du}{d\tau} - \frac{1}{2M} \left(p_u^2 + 2MV_1 u^2 - 2MV_2 + \frac{p_\phi^2}{u^2} \right) \right.$$

$$+ \left(-p_0\right) \frac{dt}{d\tau} + \frac{d\phi}{d\tau}\left(p_\phi - \sqrt{-2Mp_0}\right)\Bigg], \tag{8}$$

where $M = \frac{4m}{\alpha^2}$. Let us define $\omega = \sqrt{\frac{2V_1}{M}}$ as the frequency of the two oscillators. So that Eq. (8) can be rewritten as

$$A = \int d\tau \left[p_u \frac{du}{d\tau} - \frac{1}{2M}\left(p_u^2 + \frac{1}{2}M^2\omega^2 u^2 + \frac{p_\phi^2}{u^2} - 2MV_2\right)\right.$$

$$\left. + \left(-p_0\right) \frac{dt}{d\tau} + \frac{d\phi}{d\tau}\left(p_\phi - \sqrt{-2Mp_0}\right)\right] \tag{9}$$

We represent the position vector of the particle in the two dimensional polar coordinates as

$$\overrightarrow{\mathbf{u}} = (u_1, u_2) = \sqrt{u}\left(\cos\phi, \sin\phi\right). \tag{10}$$

Action in terms of (u_1, u_2) becomes

$$A = \int d\tau \left[p_{u_1}\frac{du_1}{d\tau} + p_{u_2}\frac{du_2}{d\tau} - \omega\left[\frac{p_{u_1}^2 + p_{u_2}^2}{2M\omega} + \frac{1}{2}M\omega(u_1^2 + u_2^2) - \frac{V_2}{\omega}\right]\right.$$

$$\left. + \left(-p_0\right)\frac{dt}{d\tau} - \sqrt{-2Mp_0}\frac{d\phi}{d\tau}\right] \tag{11}$$

Here one can define A' as an action of two oscillators system with mass M frequency ω and energy V_2. Thus we have

$$A' = \int d\tau \left\{p_{u_1}\frac{du_1}{d\tau} + p_{u_2}\frac{du_2}{d\tau} - \omega\left[\frac{p_{u_1}^2 + p_{u_2}^2}{2M\omega} + \frac{1}{2}M\omega(u_1^2 + u_2^2) - \frac{V_2}{\omega}\right]\right\}. \tag{12}$$

The term $\left[\frac{p_{u_1}^2 + p_{u_2}^2}{2M} + \frac{1}{2}M\omega^2(u_1^2 + u_2^2)\right] - V_2$ corresponds to the Hamiltonian of two oscillators.

The holomorphic coordinates are defined as

$$a = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = \frac{1}{\sqrt{2}}\begin{pmatrix} \sqrt{m\omega}u_1 + i\sqrt{\frac{1}{M\omega}}p_{u_1} \\ \sqrt{m\omega}u_2 + i\sqrt{\frac{1}{M\omega}}p_{u_2} \end{pmatrix} \tag{13}$$

and also conjugate dynamical variables as

$$a^\dagger = \left(a_1^* \quad a_2^*\right) = \frac{1}{\sqrt{2}}\left(\sqrt{m\omega}u_j - i\sqrt{\frac{1}{M\omega}}p_{u_j}\right), \qquad j = 1, 2. \tag{14}$$

The action in Eq. (12) is rewritten again in terms of the holomorphic coordinates

$$A'\left(a_b^\dagger, \tau_b; a_a, \tau_a\right) = V_2\left(\tau_b - \tau_a\right) + \int d\tau \left[\frac{1}{2i}\left(\frac{da^\dagger}{d\tau}a - a^\dagger\frac{da}{d\tau}\right) - \omega\left(a^\dagger a\right)\right]. \tag{15}$$

We ignore the total derivative $\frac{1}{2}\sum\limits_{j=1}^{2}\left(p_{u_j}u_j\right)\Big|_{\tau_a}^{\tau_b}$ in the action, Eq. (15). The kernel of two oscillators with parametric time τ in holomorphic coordinates is

$$K'(a_b^\dagger,\tau_b;a_a,\tau_a)=e^{i(V_2-)(\tau_b-\tau_a)}\int\frac{Da^\dagger Da}{[2\pi i]^2}\exp\left[i\int\limits_{\tau_a,a_a}^{\tau_b,a_b^\dagger}d\tau\frac{1}{2i}\left(\frac{da^\dagger}{d\tau}a-a^\dagger\frac{da}{d\tau}\right)-\omega\left(a^\dagger a\right)\right].\tag{16}$$

Here we take $\hbar=1$ and the last term ω is appeared from the quantum ordering terms between the operators $\widehat{a}^\dagger$ and $\widehat{a}$. After the integrations over a, $a^\dagger$ [9] we get

$$K'(a_b^\dagger,\tau_b;a_a,\tau_a)=e^{i(V_2-\omega)(\tau_b-\tau_a)}\exp\left[a_b^\dagger a_a e^{-i\omega(\tau_b-\tau_a)}\right].\tag{17}$$

Since the transformation in Eq. (16) has double value, the physical kernel becomes

$$K'(a_b^\dagger,\tau_b;a_a,\tau_a)=e^{i(V_2-\omega)(\tau_b-\tau_a)}$$
$$\times\left\{\exp\left[a_b^\dagger a_a e^{-i\omega(\tau_b-\tau_a)}\right]+\exp\left[-a_b^\dagger a_a e^{-i\omega(\tau_b-\tau_a)}\right]\right\}.\tag{18}$$

If the exponential term is expanded into power series of $a_b^\dagger$ we have

$$K'\left(a_b^\dagger,\tau_b;a_a,\tau_a\right)=e^{\tau_b-\tau_a}\sum_{n_{1,2}=0}^{\infty}\left[1+(-1)^{n_1+n_2}\right]e^{-i[(n_1+n_2+1)\omega-V_2](\tau_b-\tau_a)}\prod_{j=1}^{2}\frac{\left(a_{jb}^* a_{ja}\right)^{n_j}}{\Gamma\left(n_j+1\right)}\tag{19}$$

where n_1 and n_2 are quantum numbers. They can be expressed in terms of the radial and angular quantum numbers n_r and m of two oscillators as

$$n_1=n_r+\frac{|m|+m}{2}\ ,\ n_2=n_r+\frac{|m|-m}{2}\ .\tag{20}$$

and let us take $a_\mp$ final eigenstates

$$a_\mp=\frac{\left(a_{1b}\mp ia_{2b}\right)^*}{\sqrt{2}}\tag{21}$$

and $\lambda_\mp$ initial eigenstates

$$\lambda_\mp=\frac{\left(a_{1a}\mp ia_{2a}\right)}{\sqrt{2}}.\tag{22}$$

Thus, we rewrite the Kernel in Eq.(19) as

$$K'\left(a_\mp,\tau;\lambda_\mp;\tau_a\right)=\sum_{n_r=0}^{\infty}\sum_{m=-\infty}^{\infty}e^{-2i\left[\left(n_r+\frac{|m|}{2}+\frac{1}{2}\right)\omega-\frac{V_2}{2}\right](\tau_b-\tau_a)}$$
$$\times\left[1+(-1)^{2|m|}\frac{\left(a_+^*\lambda_+\right)^{n_1}\left(a_-^*\lambda_-\right)^{n_2}}{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}\right].\tag{23}$$

We suppose that ϕ varies as $-\pi < \phi < \pi$ for derive in Eq. (23). However the generalized Morse potential is not periodic in this interval. If we let $\phi \longrightarrow 2\pi\phi/2L$ and taking the limit $L \longrightarrow \infty$ Eq. (23), it becomes

$$K'\left(a_{\mp},\tau;\lambda_{\mp};\tau_a\right)=\sum_{n_r=0}^{\infty}\int_{-\infty}^{\infty}dm\ e^{-2i\left[\left(n_r+\frac{|m|}{2}+\frac{1}{2}\right)\omega-\frac{V_2}{2}\right]\left(\tau_b-\tau_a\right)}$$

$$\left[1+(-1)^{2|m|}\frac{\left(a_+^*\lambda_+\right)^{n_1}\left(a_-^*\lambda_-\right)^{n_2}}{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}\right]. \tag{24}$$

K' can be written in terms of oscillator energy eigenstates $|n_r,m\rangle$ as

$$K'\left(a_{\mp},\tau;\lambda_{\mp};\tau_a\right)=\sum_{n_r=0}^{\infty}\int_{-\infty}^{\infty}dm\ \ \langle n_r,m|\,a_{\mp}\rangle^*\,\langle n_r,m|\,U\left(\tau_b-\tau_a\right)|\lambda_{\mp}\rangle\,, \tag{25}$$

where $U\left(\tau_b-\tau_a\right)$ is parametric time evolution operator between initial coherent states of the two oscillators. We can denote

$$|a_{\mp}\rangle=e^{\left(\tau_b-\tau_a\right)}\sum_{n_r=0}^{\infty}\int_{-\infty}^{\infty}dm\left[1+(-1)^{|m|}\right]\frac{\left(a_+^*\right)^{n_1}\left(a_-^*\right)^{n_2}}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}} \tag{26}$$

and

$$U\left(\tau_b-\tau_a\right)|\lambda_{\mp}\rangle=\sum_{n_r=0}^{\infty}\int_{-\infty}^{\infty}dm e^{-2i\left[\left(n_r+\frac{|m|}{2}+\frac{1}{2}\right)\omega-\frac{V_2}{2}\right]\left(\tau_b-\tau_a\right)}\left[1+(-1)^{|m|}\right]\frac{\left(\lambda_+\right)^{n_1}\left(\lambda_-\right)^{n_2}}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}}. \tag{27}$$

To get the coherent states as a function of physical coordinates, Green's function of the system is obtained as

$$G'\left(u_b,t_b;\lambda_{\mp},t_a\right)=\sum_{n_r=0}^{\infty}\int_{-\infty}^{\infty}dm\frac{-i\left[1+(-1)^{2|m|}\right]}{2\omega\left(n_r+|m|+\frac{1}{2}\right)-V_2}\frac{\left(a_+^*\lambda_+\right)^{n_1}\left(a_-^*\lambda_-\right)^{n_2}}{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}. \tag{28}$$

The poles of Green's function give the energy eigenvalues of the system. In order to obtain the Green's function of the physical problem, the dummy coordinates should be eliminated. One of the methods to eradicate a variable in the path integration formalism is to integrate over it. The other method is to take the physical eigenvalues of the corresponding conjugate momenta in the wave function formalism. To eliminate the dummy coordinate ϕ, we prefer method in the wave function formalism [16]. We take m as an azimuthal quantum number corresponding to the operator $\widehat{p}_\phi$. Thus the Green's function is converted into

$$G'\left(a_\mp;\lambda_\mp\right)=\sum_{n_r=0}^{\infty}\left[1+(-1)^{|m|}\right]\frac{\left(a_+^*a_-^*\lambda_+\lambda_-\right)^{n_r+|m|}\left(\frac{a_+^*\lambda_+}{a_-^*\lambda_-}\right)^m}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}} \tag{29}$$

and physical coherent states become

$$\left|a_\mp\right\rangle=\sum_{n_r=0}^{\infty}\left[1+(-1)^{|m|}\right]\frac{\left(a_+a_-\right)^{n_r+|m|}\left(\frac{a_+}{a_-}\right)^m}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}}. \tag{30}$$

As a function of physical coordinates u and t, kernel of the system is written as

$$K'\left(u,\phi;\lambda_\mp(\tau)\right)=\int\frac{da_\mp^*da_\mp}{\left(2\pi i\right)^2}e^{-a_\mp^*a_\mp}\left\langle u,\phi\right|\left.a_\mp\right\rangle K'\left(a_\mp^*,\tau;\lambda_\mp,\tau_a\right), \tag{31}$$

with $\left\langle u,\phi\right|\left.a_\mp\right\rangle$ is

$$\left\langle u,\phi\right|\left.a_\mp\right\rangle=Ne^{-\frac{|a|^2}{2}}\exp\left[-\frac{1}{2}M\omega u^2+2\sqrt{M\omega}u\left(a_+e^{-i\phi}+a_-e^{i\phi}\right)-\frac{a^2}{2}\right]. \tag{32}$$

where $a^2=a_+a_-+a_-a_+$ and $N=\left(\frac{M\omega}{\pi}\right)^2e^{-\frac{|a|^2}{2}}$. If we take integrations over $a_\mp^*$, $a_\mp$ in Eq. (31), it will be

$$\begin{aligned}K\left(u,\phi;\lambda_\mp\right)=N\exp\Bigg[&-\frac{1}{2}M\omega u^2+2\sqrt{M\omega}ue^{-i\omega\tau}\\&\times\left(\lambda_+e^{-i\phi}+\lambda_-e^{i\phi}\right)-\frac{\lambda^2}{2}\Bigg].\end{aligned} \tag{33}$$

The parametric time dependence of the eigenvalues of the operators $\widehat{a_j}$ is the form of $a_j(\tau)=a_{ja}e^{-i\omega(\tau-\tau_a)}$. Let us rewrite it in terms of trigonometric functions

$$K\left(u,\phi,;\lambda_\mp(\tau)\right)=Ne^{-\frac{1}{2}M\omega u^2-\frac{|\lambda|^2}{2}}\exp\left[-2i\sqrt{\frac{M\omega u^2}{2}\left(i\lambda\right)^2}\left(e^{-i(\phi+\delta)}+e^{i(\phi+\delta)}\right)\right]. \tag{34}$$

Here δ is complex phase and determined as

$$e^{-i\delta}=\frac{\lambda_+-i\lambda_-}{\sqrt{\lambda_+\lambda+\lambda_-\lambda_+}}. \tag{35}$$

To get coherent states in the parabolic coordinates, we express the Kernel in terms of Bessel functions [17]

$$K\left(u,\phi,;\lambda_\mp(\tau)\right)=Ne^{-\frac{1}{2}M\omega u^2-\frac{\lambda^2}{2}}\sum_{m=-\infty}^{\infty}\left[1+(-1)^{|m|}\right].$$

$$\times J_m\left(2\sqrt{\frac{M\omega u^2}{2}\left(i\lambda\right)^2}\right)e^{-im(\phi+\delta)} \tag{36}$$

Integrating the Kernel over the parametric time τ Green's function can be written

$$G\left(u;\lambda_\mp\right)=\int\frac{dp_0}{2\pi i}e^{-ip_0(t-t_a)}\sum_{n_r=0}^{\infty}\frac{i\left[1+(-1)^{|m|}\right]}{\omega\left[n_r+|m|/2+1/2\right]-V_2/2}\;\Phi_{n_r,m}(u)$$

$$\times\frac{e^{-|\lambda|^2}e^{im\pi/2}e^{-\lambda^2/2}\left(-\lambda^2\right)^{n_r+|m|/2}}{\sqrt{\Gamma\left(n_r+|m|+1\right)}}\cos m\left(\delta+\frac{\pi}{2}\right) \tag{37}$$

where $\Phi_{n_r,m}(u)$ is

$$\Phi_{n_r,m}(u)=\frac{e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{2}M\omega/\pi\right)^{1/4}\left(\sqrt{\frac{M\omega u^2}{2}}\right)^{|m|}}{\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right). \tag{38}$$

Thus the Green's function of physical time in the parabolic coordinates can be obtained as

$$G(u,\phi,t;\lambda_\mp,t_a)=\sum_{n_r=0}^{\infty}\sum_{m=-\infty}^{\infty}e^{-iV_2\left(1-\frac{\omega}{V_2}(n_r+1/2)\right)^2(t-t_a)}\frac{e^{im\phi}}{2\pi}$$

$$\times\frac{\left(\lambda_+\lambda_-\right)^{n_r+|m|}\left(\frac{\lambda_+}{\lambda_-}\right)^m}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}}$$

$$\times\frac{\sqrt{MV_2/\left(n+1\right)}e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}(\frac{M\omega u^2}{2}). \tag{39}$$

where $n=n_r+|m|/2$. From Eq. (39) energy eigenstates become

$$\psi_{n_r,m}(u)=\sqrt{\frac{2M}{n}}\frac{\sqrt{MV_2/2n+1}^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right)$$

Using the residue of Green's function the energy eigenvalues of the system becomes

$$E=-V_2\left[1-\frac{\omega}{V_2}(n_r+1/2)\right]^2. \tag{40}$$

The above result agrees with one available in literature [13].

III. PT-SYMMETRIC AND NON-HERMITIAN GENERALIZED MORSE CASE

If V_1 and V_2 are real and $\alpha = i\alpha$ then The generalized Morse potential has the form

$$V(x) = V_1 e^{-2i\alpha x} - V_2 e^{-i\alpha x}. \tag{41}$$

We can obtain the parametric time coherent states and energy spectra for this potential following the steps of Sec. II by using the same variables then the action becomes

$$\begin{aligned} A = \int d\tau \Bigg[p_{u_1}\frac{du_1}{d\tau} + p_{u_2}\frac{du_2}{d\tau} - \omega \left[\frac{p_{u_1}^2 + p_{u_2}^2}{2M\omega} + \frac{1}{2}M\omega(u_1^2 + u_2^2) + \frac{V_2}{\omega} \right] \\ + (-p_0)\frac{dt}{d\tau} + \frac{d\phi}{d\tau}\left(p_\phi - \sqrt{2Mp_0}\right) \Bigg] \end{aligned} \tag{42}$$

Frequency of the two oscillators system is again $\omega = \sqrt{\frac{-2V_1}{M}}$ and energy is V_2. Here we see that the kinetic energy term is negative. By holomorphic coordinate transformations the action becomes

$$A'(a_b^\dagger, \tau_b; a_a, \tau_a) = \int d\tau \left[\frac{1}{2i}\left(\frac{da^\dagger}{d\tau}a - a^\dagger \frac{da}{d\tau} \right) - \omega\left(a^\dagger a\right) + V_2 \right]. \tag{43}$$

The action of the system becomes as in Eq. (15) again. The form in Eq. (16) takes

$$K'(a_b^\dagger, \tau_b; a_a, \tau_a) = e^{i(-V_2-\omega)(\tau_b-\tau_a)} \int \frac{Da^\dagger Da}{\left[2\pi i\right]^2} \exp\left\{ i \int\limits_{\tau_a, a_a}^{\tau_b, a_b^\dagger} d\tau \left[\frac{1}{2i}\left(\frac{da^\dagger}{d\tau}a - a^\dagger\frac{da}{d\tau} \right) - \omega\left(a^\dagger a\right) \right] \right\}. \tag{44}$$

So, following the same steps of Sec. II, kernel of the system is obtained as

$$K'\left(u, \phi, ; \lambda_\mp(\tau)\right) = e^{-\frac{1}{2}M\omega u^2 - \frac{\lambda_+\lambda_-}{2}} \exp\left[-2i\sqrt{\frac{M\omega u^2}{2}\left(i\lambda\right)^2}\left(e^{-i(\phi+\delta)} + e^{i(\phi+\delta)}\right) \right] \tag{45}$$

and Green's function as a function of physical coordinates is written

$$G'\left(u_b, t_b; \lambda_\mp, t_a\right) = \sum_{n_r=0}^{\infty} \int\limits_{-\infty}^{\infty} dm \frac{i\left[1 + (-1)^{2|m|}\right]}{V_2/2 - \omega\left(n_r + |m| + \frac{1}{2}\right)} \frac{\left(a_+^*\lambda_+\right)^{n_1}\left(a_-^*\lambda_-\right)^{n_2}}{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}. \tag{46}$$

Hence parametric time coherent states for PT-symmetric and non-Hermitian Generalized Morse can be written as

$$|a_\mp\rangle = \sum_{n_r=0}^{\infty} \left[1 + (-1)^{|m|}\right] \frac{\left(a_+a_-\right)^{n_r+|m|}\left(\frac{a_+}{a_-}\right)^m}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right).}} \tag{47}$$

After performing integrations again, Green's function of generalized Morse Potential takes

$$G\left(u;\lambda_{\mp}\right)=\int\frac{dp_0}{2\pi i}e^{-ip_0(t-t_a)}\sum_{n_r=0}^{\infty}\frac{i\left[1+(-1)^{|m|}\right]}{V_2/2-\omega\left[n_r+\left|m\right|/2+1/2\right]}\ \Phi_{n_r,m}(u)$$

$$\times\frac{e^{im\pi/2}e^{-\lambda^2/2}\left(-\lambda^2\right)^{n_r+|m|/2}}{\sqrt{\Gamma\left(n_r+\left|m\right|+1\right)}}\cos m\left(\delta+\frac{\pi}{2}\right), \tag{48}$$

where $\Phi_{n_r,m}(u)$ is

$$\Phi_{n_r,m}(u)=\frac{e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{\frac{M\omega u^2}{2}}\right)^{|m|}}{\sqrt{\Gamma\left(n_r+\left|m\right|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right). \tag{49}$$

Thus for PT-symmetric and non-Hermitian generalized Morse potential, Green's functions of physical time in the parabolic coordinates can be obtained

$$G(u,\phi,t;\lambda_{\mp},t_a)=\sum_{n_r=0}^{\infty}\sum_{m=-\infty}^{\infty}e^{-iV_2\left(1+\frac{\omega}{V_2}(n_r+1/2)\right)^2(t-t_a)}\frac{e^{im\phi}}{2\pi}$$

$$\times\frac{\left(\lambda_+\lambda_-\right)^{n_r+|m|}\left(\frac{\lambda_+}{\lambda_-}\right)^{m}}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}}$$

$$\times\frac{\sqrt{MV_2/2n+1}e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+\left|m\right|+1\right)}}L_{n_r+|m|}^{|m|}(\frac{M\omega u^2}{2}). \tag{50}$$

The eigenstates of PT-symmetric and non-Hermitian generalized Morse potential are

$$\psi_{n_r,m}(u)=\sqrt{\frac{2M}{n}}\frac{\sqrt{MV_2\left(n+1\right)}e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+\left|m\right|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right). \tag{51}$$

Since the energy of the system is a residue of Green's function, using the residue of Eq. (47), we get

$$E=-V_2\left[1+\frac{\omega}{V_2}(n+1/2)\right]^2. \tag{52}$$

From above equation, one can easily see that there are only real spectra for the PT-symmetric and non-hermitian Morse case [4, 7, 10, 13, 18].

IV. NON-PT-SYMMETRIC AND NON-HERMITIAN GENERALIZED MORSE CASE

If we take $V_1 = (A + iB)^2$, $V_2 = (2C + 1)(A + iB)$ and $\alpha = 1$ the generalized Morse potential can be rewritten as

$$V(x) = (A + iB)^2 e^{-2x} - (2C + 1)(A + iB) e^{-x}. \tag{53}$$

Here A, B, C are arbitrary parameters. This potential is non-PT symmetric and non-Hermitian, but it has reel spectra. If V_1 is real $V_2 = A + iB$ and $\alpha = i\alpha$ is taken then the Morse potential is transformed into the form

$$V(x) = V_1 e^{-2i\alpha x} - (A + iB) e^{-i\alpha x}. \tag{54}$$

Now we can derive coherent states and energy spectra following the same steps in part II we get the action as

$$A = \int ds \left\{ (-p_0) \frac{dt}{d\tau} + \sqrt{2mp_0} \frac{d\phi}{d\tau} + A' \right\} \tag{55}$$

where ϕ is a dummy coordinate. Lagrange multiplier $\sqrt{2mp_o}$ is added. Hence the new action becomes

$$A' = \int d\tau \left\{ p_{u_1} \frac{du_1}{d\tau} + p_{u_2} \frac{du_2}{d\tau} - \omega \left[\frac{p_{u_1}^2 + p_{u_2}^2}{2M\omega} + \frac{1}{2} M\omega \left(u_1^2 + u_2^2\right) - \frac{A + iB}{\omega} \right] \right\} \tag{56}$$

Here A' is two oscillators action and frequency $\omega = \sqrt{\frac{-V_1}{M}}$ and energy is $A + iB$. Here we notice that the kinetic energy term is negative. The action can be written by holomorphic coordinate transformation as

$$A'(a_b^\dagger, \tau_b; a_a, \tau_a) = \int d\tau \left[\frac{1}{2i} \left(\frac{da^\dagger}{d\tau} a - a^\dagger \frac{da}{d\tau} \right) - \omega \left(a^\dagger a\right) - (A + iB) \right]. \tag{57}$$

The kernel of the system becomes

$$K'(a_b^\dagger, \tau_b; a_a, \tau_a) = \int \frac{Da^\dagger Da}{[2\pi i]^2} \exp \left\{ i \int d\tau \left[\frac{1}{2i} \left(\frac{da^\dagger}{d\tau} a - a^\dagger \frac{da}{d\tau} \right) \right.\right.$$
$$\left.\left. - \omega \left(a^\dagger a\right) - (A + iB) \right] \right\}. \tag{58}$$

So, Green's function of the physical system can be obtained as follows

$$G'(u_b, t_b; \lambda_\mp, t_a) = \sum_{n_r=0}^{\infty} \int_{-\infty}^{\infty} dm \frac{i \left[1 + (-1)^{2|m|}\right]}{A + iB - \omega \left(n_r + |m| + \frac{1}{2}\right)} \frac{\left(a_+^* \lambda_+\right)^{n_1} \left(a_-^* \lambda_-\right)^{n_2}}{\Gamma(n_1 + 1)\Gamma(n_2 + 1)} \tag{59}$$

and the parametric time coherent states for non-PT-symmetric and non-Hermitian generalized Morse potential become are the same with the ones given in Eq. (30). After performing integrations, wave functions of the coherent states are

$$G\left(u;\lambda_{\mp}\right)=\int\frac{dp_0}{2\pi i}e^{-ip_0(t-t_a)}\sum_{n_r=0}^{\infty}\frac{i\left[1+(-1)^{|m|}\right]}{A+iB-\omega\left[n_r+|m|/2+1/2\right]}\,\Phi_{n_r,m}(u)$$

$$\times\frac{e^{im\pi/2}e^{-\lambda^2/2}\left(-\lambda^2\right)^{n_r+|m|/2}}{\sqrt{\Gamma\left(n_r+|m|+1\right)}}\cos m\left(\delta+\frac{\pi}{2}\right), \tag{60}$$

where $\Phi_{n_r,m}(u)$ is

$$\Phi_{n_r,m}(u)=\frac{e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{\frac{M\omega u^2}{2}}\right)^{|m|}}{\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right). \tag{61}$$

Therefore for non-PT-symmetric and non-Hermitian Morse potential, the wave function of physical time in the parabolic coordinates can be obtained as

$$G(u,\phi,t;\lambda_{\mp},t_a)=\sum_{n_r=0}^{\infty}\sum_{m=-\infty}^{\infty}e^{-i(A+iB)\left(1+\frac{\omega}{A+iB}(n_r+1/2)\right)^2(t-t_a)}\frac{e^{im\phi}}{2\pi}$$

$$\times\frac{\left(\lambda_+\lambda_-\right)^{n_r+|m|}\left(\frac{\lambda_+}{\lambda_-}\right)^m}{\sqrt{\Gamma\left(n_1+1\right)\Gamma\left(n_2+1\right)}}$$

$$\times\frac{\sqrt{M\left(A+iB\right)\left(n+1\right)}e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}(\frac{M\omega u^2}{2}). \tag{62}$$

The eigenstates of non-PT-symmetric and non-Hermitian generalized Morse case are

$$\psi_{n_r,m}(u)=\sqrt{\frac{2M}{n}}\frac{\sqrt{M\left(A+iB\right)n+1}e^{-\frac{1}{2}M\omega u^2}\left(\sqrt{M\omega u^2/2}\right)^{|m|/2}}{\pi^{1/4}\sqrt{\Gamma\left(n_r+|m|+1\right)}}L_{n_r+|m|}^{|m|}\left(\frac{M\omega u^2}{2}\right). \tag{63}$$

Since the energy of the system is a residue of Green's function, from the residue of Eq. (58), we obtain energy eigenvalues as

$$E=-(A+iB)\left[1+\frac{\omega}{A+iB}(n+1/2)\right]^2. \tag{64}$$

It is clear that energy spectra are real in cases $V_1>0$ if and only if $Re(V_2)=0$ and $V_1<0$ if and only if $Im(V_2)=0$.

V. CONCLUSION

We have studied parametric time coherent states for PT-/non-PT-Symmetric and non-Hermitian generalized Morse potential by using path integral formalism. Energy eigenvalues and the corresponding wave functions are calculated. We have discussed the negative energy coherent states. The wave functions of the potential in Secs. III and IV are physical like Sec. II. Energy eigenvalues are positive in contrary to expectation. Negative energy coherent states can be obtained by analytic continuation like [9].

VI. ACKNOWLEDGEMENTS

This research was partially supported by the Scientific and Technological Research Council of Turkey.

---

[1] C. M. Bender and S.Boettcher, Pys. Lett., **80**, 5243 (1998); C. M. Bender, S. Boetcher, and P.N. Meisenger, J. Math. Phys. **40**, 2201 (1999); C. M. Bender, G. V. Dunne, and P. N. Meisenger, Phys. Lett. **A252**, 272 (1999).

[2] Z.Ahmed, Phys. Lett. **A282**, 343 (2001).

[3] B. Bagchi, C. Quesne, Phys. Lett. **A273**, 285 (2000).

[4] C. Berkdemir, A. Berkdemir and R.Sever, Physical Rev. **C72**,027001 (2005).

[5] M. Znojil, Phys. Lett. **A264**, 108 (1999).

[6] R.P.Feynman, Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill, New York, (1965). chap. 1-2

[7] N. Kandirmaz, R. Sever, Chin. J. Phys. **47**, 46 (1999) [`arXiv:0812.2614`].

[8] P. M. Morse. Phys. Rev. **34**, 57 (1929).

[9] N. Unal, Phys. Can. J. Phys. **80**, 875 (2002).

[10] C. Berkdemir, J. Han, Chem. Phys.Lett. **409**, 203 (2005).

[11] I.H. Duru, Phys. Rev. **D28**, 10 (1983).

[12] I.H. Duru and H. Kleinert, Phys. Lett. **B84**, 185 (1979) Fortschr. Phys. **30**, 401 (1982).

[13] M. Aktas, R. Sever, Journal of Molecular Structure-Theochem **710**, 223 (2004).

[14] E. Schrödinger, Naturwissenschaften, **14**, 664 (1926).

[15] N. Unal, Found Phys. **28**, 755 (1998).

[16] N. Kandirmaz, N. Unal, Theor. and Math. Phys. **155**, 884 (2008).

[17] I.S. Gradshteyn and I.M. Ryzhic:Table of Integrals, Series, and Products,2nd ed. Academic Press, New York, (1981), chap.8.

[18] O. Yesiltas, M. Simsek, R. Sever and C. Tezcan, Phys. Script. **67**, 472 (2003).